

National Library
of Medicine

PubMed

PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM
Search PubMed	▼ for					Go	Clear
Limits Preview/Index History Clipboard							

Entrez PubMed

☐ 1: *Dev Biol Stand* 1994;82:215-27[Related Articles, Books, LinkOut](#)PubMed
Services**Mucosal vaccines based on the use of cholera toxin B subunit as immunogen and antigen carrier.****Lebens M, Holmgren J**

Department of Medical Microbiology and Immunology, University of Goteborg, Sweden.

Related
Resources

Stimulation of strong mucosal IgA immune responses as a basis for vaccine-induced protection against various pathogens has proved difficult. Most soluble protein antigens administered either parenterally or oral-mucosally have given disappointing results. A notable exception in this regard are cholera toxin (CT) and, particularly in humans, its non-toxic B subunit pentamer moiety (CTB) both of which stimulate a strong intestinal IgA antibody response and long-lasting immunological memory. Based on this, CTB has become an important component in recently developed oral vaccines against cholera and diarrhea caused by enterotoxigenic *E. coli*. The strong immunogenicity of CT and CTB can to a large extent be explained by their ability to bind to receptors on the intestinal mucosal surface. This has promoted much recent interest in the use of CTB as an oral delivery carrier for other vaccine-relevant antigens. Oral administration of antigens coupled to CTB either chemically or genetically has in several systems been found to markedly potentiate both intestinal and extra-intestinal IgA immune responses against the CTB-coupled antigens and to elicit substantial circulating antibody responses. In contrast to CTB, CT also has strong adjuvant properties for stimulating mucosal IgA immune responses to unrelated, non-coupled antigens after oral co-immunization. This adjuvant activity appears to be closely linked to the A subunit-catalyzed ADP-ribosylating action of CT leading to enhanced cyclic AMP formation in the affected cells.

Publication Types:

- Review
- Review, tutorial

PMID: 7958476, UI: 95046935

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)

is la Abstract a e e t Order Add to li oard

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)

affected cell, and efforts to eliminate the enterotoxigenic activity without losing adjuvant activity have so far not met with success.

Publication Types:

- Review
- Review, tutorial

PMID: 8203723, UI: 94263008

Display	Abstract	<input type="button" value="v"/>	Save	Text	Order	Add to Clipboard
---------	----------	----------------------------------	------	------	-------	------------------

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)

National Library
of Medicine

PubMed

PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM
Search PubMed	for					Go	Clear
Limits		Preview/Index		History		Clipboard	

Entrez PubMed

☐ 1: *Int Arch Allergy Immunol* 1999
Feb-Apr; 118(2-4):129-32

Related Articles, Books,
LinkOut

KARGER
Full TextPubMed
Services

Modulation of an allergic immune response via the mucosal route in a murine model of inhalative type-I allergy.

Wiedermann U, Jahn-Schmid B, Repa A, Kraft D, Ebner C

Institute of General and Experimental Pathology, Division Immunopathology,
University of Vienna, Austria. ursula.wiedermann@akh-wien.ac.at

Related
Resources

A murine model of aerosol inhalation, leading to sensitization to birch pollen (BP) and its major allergen Bet v 1, was established in order to try to influence type-I allergic immune responses via the mucosal route. We previously demonstrated that simultaneous inhalation of BP and cholera toxin, a potent mucosal adjuvant, induced a Th1-like immune response to the allergen in naive mice and modulated allergic immune responses in sensitized mice. In contrast to cholera holotoxin, mucosal application of the cholera B subunit (CTB) conjugated to antigen has been shown to induce peripheral tolerance in certain models of Th1-based autoimmune diseases. In the present study we investigated the potential of such an antigen delivery system to suppress Th2-based, allergic immune responses. Mucosal administration of CTB/Bet v 1 conjugates prior to sensitization led to significantly increased allergen-specific IgE/IgG1 and IgG2a antibody levels and cytokine production (IL-5, IFN-gamma) in vitro. Thus, CTB coupled to Bet v 1 acted as an adjuvant rather than a tolerogen. On the other hand we noted that mucosal application of CTB coupled to ovalbumin led to marked suppression of antigen-specific IgE antibody levels and IL-5 production in vitro and thereby restricted allergic sensitization. These results indicated that the effects of CTB/antigen conjugates depended on the nature of the antigen. In contrast to Bet v 1 coupled to CTB, nasal as well as oral application of low doses of unconjugated, Bet v 1 prior to aerosol sensitization inhibited allergen-specific antibody responses of all isotypes, cutaneous type-I skin tests in vivo as well as allergen-specific lymphoproliferative responses and cytokine production (IL-4, IL-5, IL-10, IFN-gamma) in vitro, suggesting that both T- and B-cell tolerance to the allergen were induced. Taken together, mucosal tolerance induction as well as the use of certain transmucosal antigen delivery systems might be promising new strategies to modulate type-I allergic immune responses

PMID: 10224359, UI: 99242414

National Library
of Medicine

PubMed

PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM
Search PubMed	▼ for					Go	Clear
Limits		Preview/Index		History		Clipboard	

Entrez PubMed

☐ 1: *Int Immunol* 1999 Jul;11(7):1131-8

Related Articles, Books, LinkOut

[Full text article at
intimm.oupjournals.org](http://intimm.oupjournals.org)PubMed
ServicesCorrected and republished in *Int Immunol* 1999 Oct;11(10):1717-24**Suppressive versus stimulatory effects of allergen/cholera toxoid (CTB) conjugates depending on the nature of the allergen in a murine model of type I allergy.****Wiedermann U, Jahn-Schmid B, Lindblad M, Rask C, Holmgren J, Kraft D, Ebner C**Related
Resources

Division of Immunopathology, Institute of General and Experimental Pathology, University of Vienna, Waehringer Guertel 18-20, 1090 Vienna, Austria.

Recent reports have demonstrated that feeding small amounts of antigen conjugated to the B subunit of cholera toxin (CTB) suppress immune responses in experimental models of certain Th1-based autoimmune diseases. We have established a model of aerosol sensitization leading to Th2-mediated allergic immune responses in BALB/c mice. In the present study two different antigens, the dietary antigen ovalbumin (OVA) and the inhalant allergen Bet v 1 (the major birch pollen allergen), chemically coupled to recombinant CTB were tested for their potential to influence Th2-like immune responses. Intranasal administration of OVA-CTB prior to sensitization with OVA led to a significant decrease of antigen-specific IgE antibody levels, but a marked increase of OVA-specific IgG2a antibodies as compared to non-pretreated, sensitized animals. Antigen-specific lympho-proliferative responses in vitro were reduced by 65% in the pretreated group; IL-5 and IL-4, but not IFN-gamma, production were markedly decreased in responder cells of lungs and spleens of nasally pretreated mice. In contrast, mucosal administration of rBet v 1-CTB conjugates prior to sensitization led to an up-regulation of allergen-specific IgE, IgG1 and IgG2a, increased in vitro lympho-proliferative responses as well as augmented production of IL-5, IL-4, IL-10 and IFN-gamma. Intranasal administration prior to sensitization of unconjugated allergens showed also contrasting effects: OVA could not significantly influence antigen-specific antibody or cytokine production, whereas intranasal pretreatment with unconjugated Bet v 1 suppressed allergen-specific immune responses in vivo and in vitro. These results demonstrated that the two

Reviewed

National Library
of Medicine

PubMed

PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM
Search PubMed	▼ for					Go	Clear
Limits		Preview/Index		History		Clipboard	

Display	Abstract	▼	Save	Text	Order	Add to Clipboard
---------	----------	---	------	------	-------	------------------

Entrez PubMed


☐ 1: *Am J Trop Med Hyg* 1994;50(5 Suppl):42-54[Related Articles, Books, LinkOut](#)PubMed
Services**Strategies for the induction of immune responses at mucosal surfaces making use of cholera toxin B subunit as immunogen, carrier, and adjuvant.****Holmgren J, Czerkinsky C, Lycke N, Svennerholm AM**

Department of Medical Microbiology and Immunology, University of Goteborg, Sweden.

Related
Resources

The concept of a common mucosal immune system, through which specific antigen-activated lymphocytes from the gut can disseminate immunity both along the intestinal tract and to various other mucosal and glandular tissues, has generated much current interest in the possibility of developing oral vaccines, not only for enteric infections but also for infections in the respiratory and urogenital tracts. However, to date it has proven difficult in practice to stimulate strong mucosal IgA immune responses by either parenteral or oral-mucosal administration of most antigens, and experience with soluble protein antigens has, on the whole, been disappointing. A notable exception in this regard is cholera toxin (CT) and in humans more than in other species, its nontoxic B subunit pentamer moiety (CTB). Based on this, CTB has become an important component in recently developed oral vaccines against cholera as well as against diarrhea caused by enterotoxigenic *Escherichia coli* producing CT-like heat-labile enterotoxin(s). Since the strong immunogenicity of CT and CTB can, to a large extent, be explained by their ability to bind to receptors on the intestinal mucosal surface, there has recently been much interest in approaches using CTB as an oral delivery carrier system for other vaccine-relevant antigens, and much progress has been made in preparing immunogenic hybrid proteins by coupling various protein or peptide antigens chemically or genetically to CTB. Indeed, in several systems, oral administration of such hybrid antigens has been found to markedly potentiate both intestinal and extraintestinal IgA immune responses against the CTB-coupled antigens and also to elicit substantial circulating antibody responses. Besides the mucosal immunopotentiating effect of either CT or CTB owing to their similar capacity as oral antigen-delivery vehicles, CT, but in most systems tested not CTB, also has strong adjuvant properties for stimulating mucosal IgA immune responses to admixed (not coupled) unrelated antigens after oral immunization. This adjuvant activity appears to be closely linked to the ADP-ribosylating action of CT (and specifically of its A subunit) leading to enhanced cyclic AMP formation in the

Requested

National Library
of Medicine

PubMed

PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM
Search PubMed	for					Go	Clear
Limits		Preview/Index		History		Clipboard	

Display	Abstract	Save	Text	Order	Add to Clipboard
---------	----------	------	------	-------	------------------

Entrez PubMed

☐ 1: *Immunology* 1998 Jul;94(3):424-30[Related Articles, Books, LinkOut](#)**Online****Role of GM1 binding in the mucosal immunogenicity and adjuvant activity of the Escherichia coli heat-labile enterotoxin and its B subunit.**PubMed
Services**de Haan L, Verweij WR, Feil IK, Holtrop M, Hol WG, Agsteribbe E, Wilschut J**

Department of Physiological Chemistry, Groningen Utrecht Institute for Drug Exploration (GUIDE), University of Groningen, The Netherlands.

Related
Resources

Escherichia coli (E. coli) heat-labile toxin (LT) is a potent mucosal immunogen and immunoadjuvant towards co-administered antigens. LT is composed of one copy of the A subunit, which has ADP-ribosylation activity, and a homopentamer of B subunits, which has affinity for the toxin receptor, the ganglioside GM1. Both the ADP-ribosylation activity of LTA and GM1 binding of LTB have been proposed to be involved in immune stimulation. We investigated the roles of these activities in the immunogenicity of recombinant LT or LTB upon intranasal immunization of mice using LT/LTB mutants, lacking either ADP-ribosylation activity, GM1-binding affinity, or both. Likewise, the adjuvant properties of these LT/LTB variants towards influenza virus subunit antigen were investigated. With respect to the immunogenicity of LT and LTB, we found that GM1-binding activity is essential for effective induction of anti-LTB antibodies. On the other hand, an LT mutant lacking ADP-ribosylation activity retained the immunogenic properties of the native toxin, indicating that ADP ribosylation is not critically involved. Whereas adjuvanticity of LTB was found to be directly related to GM1-binding activity, adjuvanticity of LT was found to be independent of GM1-binding affinity. Moreover, a mutant lacking both GM1-binding and ADP-ribosylation activity, also retained adjuvanticity. These results demonstrate that neither ADP-ribosylation activity nor GM1 binding are essential for adjuvanticity of LT, and suggest an ADP-ribosylation-independent adjuvant effect of the A subunit.

PMID: 9767427, UI: 98444394

Display	Abstract	Save	Text	Order	Add to Clipboard
---------	----------	------	------	-------	------------------

National Library
of Medicine

PubMed

PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM
Search PubMed	▼ for					Go	Clear
Limits		Preview/Index		History	Clipboard		

Display	Abstract	▼	Save	Text	Order	Add to Clipboard
---------	----------	---	------	------	-------	------------------

Entrez PubMed

☐ 1: *Int Arch Allergy Appl Immunol* 1983;72(2):119-27[Related Articles, Books](#)PubMed
Services**IgA isotype restriction in the mucosal but not in the extramucosal immune response after oral immunizations with cholera toxin or cholera B subunit.**

Lycke N, Lindholm L, Holmgren J

Related
Resources

Intestinal mucosal as well as extramucosal antibody responses were studied in mice after peroral immunizations with cholera toxin or cholera B subunit. The immunizations with cholera toxin gave rise to a marked response with antitoxin-secreting cells (PFC) in Peyer's patches (PP), mesenteric lymph nodes (MLN) and spleen showing isotype distribution of IgG greater than IgA greater than IgM and with PFC kinetics in MLN and spleen that suggested migration of cells from PP after peroral administration rather than cells stimulated in situ by adsorbed antigen. Highest numbers of PFC were obtained after 2 immunizations, and further administrations resulted in a decrease in the PFC response in MLN and spleen, while the PP responsiveness was relatively unchanged, and interestingly, protective immunity and IgA-dominated antitoxin titers in intestinal washings increased markedly by the additional boosters. Animals immunized with cholera B subunit, which lacks the adenylate cyclase-stimulating capacity of cholera toxin, showed similar PFC responses in extramucosal organs as those receiving cholera toxin but were poorly protected and had correspondingly lower IgA antitoxin titers in intestinal washings. These results suggest that the mucosal IgA antitoxin predominance is mainly due to regulatory mechanisms operating on the end-stage differentiation of the committed B cells in lamina propria and that this differentiation, as judged from the different results with cholera toxin and its B subunit, might be influenced by cyclic AMP.

PMID: 6874106, UI: 83264489

Display	Abstract	▼	Save	Text	Order	Add to Clipboard
---------	----------	---	------	------	-------	------------------

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
Department of Health & Human Services
[Freedom of Information Act](#) | [Disclaimer](#)

National Library
of Medicine

PubMed

PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM
Search PubMed	for					Go	Clear
Limits		Preview/Index		History		Clipboard	

Entrez PubMed

☐ 1: *Vaccine* 1998 Jan-Feb;16(2-3):286-92[Related Articles, Books, LinkOut](#)PubMed
Services**Induction of mucosal and systemic immune responses by intranasal immunization using recombinant cholera toxin B subunit as an adjuvant.****Wu HY, Russell MW**

Department of Microbiology, University of Alabama at Birmingham 35294-2170, USA. medm115@uabdpdpo.uab.edu

Related
Resources

Intranasal (i.n.) immunization with *Streptococcus mutans* surface protein AgI/II mixed with cholera toxin B subunit (CTB) containing a trace amount of cholera toxin (CT) induces strong immune responses in mucosal and systemic sites, but whether pure CTB alone has an adjuvant effect has been questioned. To determine the adjuvant effect of recombinant (r) CTB, mice were immunized with 10 micrograms of AgI/II either mixed with or conjugated to 5 micrograms of rCTB, and antibody responses in saliva, nasal wash, gut wash, vaginal wash, and serum were assayed by ELISA. The results showed that AgI/II either mixed with or conjugated to rCTB could induce both mucosal IgA and systemic IgG antibodies to higher levels than in mice similarly immunized with AgI/II alone. Some responses, especially serum IgG antibodies, were enhanced by adding 5 micrograms CT to the immunogen, whereas overall mice immunized with AgI/II mixed with CTB contaminated with CT tended to generate the strongest mucosal IgA and serum IgG responses to AgI/II. However, rCTB used as an adjuvant induced lower antibody responses against itself than CTB intentionally or inadvertently mixed with CT. These results show that rCTB can serve as an adjuvant for protein immunogens administered by the i.n. route.

PMID: 9607044, UI: 98269927

Display	Abstract	Save	Text	Order	Add to Clipboard
---------	----------	------	------	-------	------------------

[Write to the Help Desk](#)[NCBI](#) | [NLM](#) | [NIH](#)[Department of Health & Human Services](#)[Freedom of Information Act](#) | [Disclaimer](#)

National Library
of Medicine

PubMed

PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM
Search PubMed	<input type="text" value="for"/>					Go	Clear
Limits Preview/Index History Clipboard							

Display	Abstract	<input type="text" value=""/>	Save	Text	Order	Add to Clipboard
---------	----------	-------------------------------	------	------	-------	------------------

Entrez PubMed

☐ 1: *Microbiol Immunol* 2000;44(4):259-66[Related Articles, Books, LinkOut](#)PubMed
Services

Intranasal sensitization of Japanese cedar pollen by the co-administration of low doses of cholera toxin but not its recombinant B subunit to mice.

Hirai T, Hashiguchi S, Torigoe N, Toda Y, Ito Y, Sugimur K


Department of Bioengineering, Faculty of Engineering, Kagoshima University, Kagoshima, Japan.

Related
Resources

We evaluated the effects of cholera toxin (CT) and the B subunit of cholera toxin (CTB) on the intranasal sensitization of Japanese cedar pollen (JCP) in mice. JCP suspended in phosphate-buffered saline was administered into the nostrils of mice in combination with varying doses of CT or recombinant CTB(r-CTB) once a week for 5 weeks. Antibody responses specific to sugi basic protein (SBP) were monitored by ELISA for seven weeks. The sensitization of JCP alone did not induce IgG1, IgG2b, IgG2a, IgE or IgA. In contrast, sensitization of JCP in combination with CT (JCP/CT) elicited the prominent production of SBP-specific IgG1 and low levels of IgG2b and IgG2a on Day 49. IgE production was detected only in the serum of mice which were treated with JCP/CT, and not under any other protocol. Using spleen cells from these mice, cytokine production was examined by ELISA in culture supernatants after they had been stimulated in vitro with major cedar pollen allergens, Cry j 1, Cry j 2 or SBP. Notable responses were an increase of IFN-gamma as well as IL-4 in JCP/CT-sensitized cells stimulated with Cry j 2, but not in those stimulated with Cry j 1. No significant differences were detected in IL-5 production among the experimental groups. Histopathological examination, however, showed that eosinophil infiltration was evident in the nasal mucosa of the JCP/CT-sensitized mice following challenge with JCP/CT, but weak with BSA/CT or CT alone. Thus, the immunological and histological analyses indicated that the co-administration of a low dose of CT in combination with JCP allows the induction of pollen-allergic states in mice.

PMID: 10832969, UI: 20290566

Display	Abstract	<input type="text" value=""/>	Save	Text	Order	Add to Clipboard
---------	----------	-------------------------------	------	------	-------	------------------

National Library
of Medicine

PubMed

PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM
Search PubMed	▼ for					Go	Clear
Limits		Preview/Index		History		Clipboard	

Entrez PubMed

Display	Summary	▼	Save	Text	Order	Add to Clipboard
Show: 20	▼	Items 1-20 of 228	Page 1 of 12		Select page: 1 2 3 4 5 6 7 8 9 10 >>	

PubMed
Services☐ 1: McKenzie SJ, Halsey JF.

Related Articles

Cholera toxin B subunit as a carrier protein to stimulate a mucosal immune response.

J Immunol. 1984 Oct;133(4):1818-24.

PMID: 6470484; UI: 84290652

☐ 2: Sugita-Konishi Y, Kumagai S.

Related Articles

Biliary immune response to orally presented food antigen, ovomucoid, and its potentiation by cholera toxin B subunit.

Scand J Immunol. 1992 May;35(5):597-602.

PMID: 1579860; UI: 92254026

Related
Resources☐ 3: Wu HY, Russell MW.

Related Articles

Comparison of systemic and mucosal priming for mucosal immune responses to a bacterial protein antigen given with or coupled to cholera toxin (CT) B subunit, and effects of pre-existing anti-CT immunity.

Vaccine. 1994;12(3):215-22.

PMID: 8165853; UI: 94219163

☐ 4: Holmgren J, Czerkinsky C, Lycke N, Svennerholm AM.

Related Articles

Strategies for the induction of immune responses at mucosal surfaces making use of cholera toxin B subunit as immunogen, carrier, and adjuvant.

Am J Trop Med Hyg. 1994;50(5 Suppl):42-54. Review.

PMID: 8203723; UI: 94263008

☐ 5: Bergquist C, Lagergard T, Lindblad M, Holmgren J.

Related Articles

Local and systemic antibody responses to dextran-cholera toxin B subunit conjugates.

Infect Immun. 1995 May;63(5):2021-5.

PMID: 7537252; UI: 95247296

☐ 6: Lebens M, Holmgren J.

Related Articles

Mucosal vaccines based on the use of cholera toxin B subunit as immunogen and antigen carrier.

Dev Biol Stand. 1994;82:215-27. Review.

PMID: 7958476; UI: 95046935

☐ 7: Wu HY, Russell MW.

Related Articles

Induction of mucosal and systemic immune responses by intranasal immunization

using recombinant cholera toxin B subunit as an adjuvant.
Vaccine. 1998 Jan-Feb;16(2-3):286-92.
PMID: 9607044; UI: 98269927

☐ 8: Stok W, van der Heijden PJ, Bianchi AT.

Related Articles

Conversion of orally induced suppression of the mucosal immune response to ovalbumin into stimulation by conjugating ovalbumin to cholera toxin or its B subunit.
Vaccine. 1994 May;12(6):521-6.
PMID: 8036826; UI: 94310831

☐ 9: Russell MW, Wu HY.

Related Articles

Distribution, persistence, and recall of serum and salivary antibody responses to peroral immunization with protein antigen I/II of Streptococcus mutans coupled to the cholera toxin B subunit.
Infect Immun. 1991 Nov;59(11):4061-70.
PMID: 1937766; UI: 92040078

☐ 10: de Geus B, Dol-Bosman M, Scholten JW, Stok W, Bianchi A.

Related Articles

A comparison of natural and recombinant cholera toxin B subunit as stimulatory factors in intranasal immunization.
Vaccine. 1997 Jul;15(10):1110-3.
PMID: 9269054; UI: 97414197

☐ 11: Bianchi AT, Zwart RJ, Van der Heijden PJ.

Related Articles

Induction of an enteric Ig-response against ovalbumin and stimulation of the response by cholera toxin and its B-subunit in mice.
Reg Immunol. 1990-91;3(3):131-8.
PMID: 2098068; UI: 91274179

☐ 12: Harokopakis E, Childers NK, Michalek SM, Zhang SS, Tomasi M.

Related Articles

Conjugation of cholera toxin or its B subunit to liposomes for targeted delivery of antigens.
J Immunol Methods. 1995 Sep 11;185(1):31-42.
PMID: 7665899; UI: 95395274

☐ 13: Menge AC, Michalek SM, Russell MW, Mestecky J.

Related Articles

Immune response of the female rat genital tract after oral and local immunization with keyhole limpet hemocyanin conjugated to the cholera toxin B subunit.
Infect Immun. 1993 May;61(5):2162-71.
PMID: 8478106; UI: 93239330

☐ 14: Wu HY, Russell MW.

Related Articles

Induction of mucosal immunity by intranasal application of a streptococcal surface protein antigen with the cholera toxin B subunit.
Infect Immun. 1993 Jan;61(1):314-22.
PMID: 8418053; UI: 93114890

☐ 15: Bergquist C, Lagergard T, Holmgren J.

Related Articles

Antibody responses in serum and lung to intranasal immunization with Haemophilus influenzae type b polysaccharide conjugated to cholera toxin B

subunit and tetanus toxoid.
APMIS. 1998 Aug;106(8):800-6.
PMID: 9744766; UI: 98416015

- ☐ 16: [de Fijter JW, Eijgenraam JW, Braam CA, Holmgren J, Daha MR, van Es LA, van den Wall Bake AW.](#) [Related Articles](#)

Deficient IgA1 immune response to nasal cholera toxin subunit B in primary IgA nephropathy.
Kidney Int. 1996 Sep;50(3):952-61.
PMID: 8872971; UI: 97026793

- ☐ 17: [Muller CP, Beauverger P, Schneider F, Jung G, Brons NH.](#) [Related Articles](#)

Cholera toxin B stimulates systemic neutralizing antibodies after intranasal co-immunization with measles virus.
J Gen Virol. 1995 Jun;76 (Pt 6):1371-80.
PMID: 7782766; UI: 95302033

- ☐ 18: [Tamura S, Funato H, Nagamine T, Aizawa C, Kurata T.](#) [Related Articles](#)

Effectiveness of cholera toxin B subunit as an adjuvant for nasal influenza vaccination despite pre-existing immunity to CTB.
Vaccine. 1989 Dec;7(6):503-5.
PMID: 2609726; UI: 90118301

- ☐ 19: [Lycke N, Lindholm L, Holmgren J.](#) [Related Articles](#)

IgA isotype restriction in the mucosal but not in the extramucosal immune response after oral immunizations with cholera toxin or cholera B subunit.
Int Arch Allergy Appl Immunol. 1983;72(2):119-27.
PMID: 6874106; UI: 83264489

- ☐ 20: [Van der Heijden PJ, Bianchi AT, Dol M, Pals JW, Stok W, Bokhout BA.](#) [Related Articles](#)

Manipulation of intestinal immune responses against ovalbumin by cholera toxin and its B subunit in mice.
Immunology. 1991 Jan;72(1):89-93.
PMID: 1705237; UI: 91147150

Display	Summary	▼	Save	Text	Order	Add to Clipboard
Show: 20	▼	Items 1-20 of 228	Page 1 of 12	Select page	1 2 3 4 5 6 7 8 9 10 >>	

[Write to the Help Desk](#)
[NCBI | NLM | NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act | Disclaimer](#)

National Library
of Medicine

PubMed

PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM
Search PubMed	for					Go	Clear
Limits		Preview/Index		History		Clipboard	

Entrez PubMed

☐ 1: *Vaccine* 1994;12(3):215-22[Related Articles, Books, LinkOut](#)PubMed
Services

Comparison of systemic and mucosal priming for mucosal immune responses to a bacterial protein antigen given with or coupled to cholera toxin (CT) B subunit, and effects of pre-existing anti-CT immunity.

Wu HY, Russell MW

Department of Microbiology, University of Alabama at Birmingham 35294.

Related
Resources

Intraperitoneal immunization with a bacterial protein antigen conjugated to cholera toxin B subunit (CTB) was generally less effective than intragastric or intranasal immunization in generating mucosal IgA antibodies, and in priming the mucosal immune system to respond to intragastric or intranasal boosting. Previous intragastric or intranasal immunization which generated high levels of mucosal and circulating antibodies to CTB did not suppress mucosal IgA responses induced by intragastric or intranasal immunization with bacterial antigen conjugated to or mixed with CTB, but serum antibody responses were inhibited depending on the route of immunization and whether CTB was conjugated to or mixed with the antigen.

PMID: 8165853, UI: 94219163

Display	Abstract	Save	Text	Order	Add to Clipboard
---------	----------	------	------	-------	------------------

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)

National Library
of Medicine

PubMed

PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM
Search PubMed	for					Go	Clear
Limits		Preview/Index		History	Clipboard		

Display	Abstract	Save	Text	Order	Add to Clipboard
---------	----------	------	------	-------	------------------

Entrez PubMed

☐ 1: *Proc Natl Acad Sci U S A* 1994 Nov
8;91(23):10795-9

Related Articles, Books,
LinkOut

FREE full text article at
www.pnas.org

PubMed
Services

Cholera toxin B subunit: an efficient transmucosal carrier-delivery system for induction of peripheral immunological tolerance.

Sun JB, Holmgren J, Czerkinsky C

Department of Medical Microbiology and Immunology, University of Goteborg, Sweden.

Related
Resources

Oral administration of antigens, including allergens and autoantigens, may be an efficient way to prevent diseases associated with untoward immune responses to self- and non-self-antigens. However, this approach has met with limitations because it usually requires repeated administrations of large doses of antigen and is less efficient in an already immune host, and the effect is of short duration. We report that a single oral administration of minute amounts of particulate or soluble antigen coupled to the B subunit of cholera toxin (CTB) can markedly suppress systemic immune responses in naive and in systemically immune animals. Both early (2-4 hr) and late (24-48 hr) delayed type-hypersensitivity reactivities were strongly suppressed after feeding a single dose of CTB-conjugated antigen. Serum antibody responses were also decreased, although moderately, after oral administration of CTB-conjugated antigen. This strategy of tolerance induction, based on oral administration of small amounts of antigens conjugated to a mucosa-binding molecule, may find broad applications for preventing or abrogating untoward immune responses.

Comments:

- Comment in: *Proc Natl Acad Sci U S A* 1994 Nov 8;91(23):10762-5

PMID: 7526379, UI: 95062151

Display	Abstract	Save	Text	Order	Add to Clipboard
---------	----------	------	------	-------	------------------

Write to the Help Desk
NCBI | NLM | NIH

National Library
of Medicine

PubMed

PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM
Search PubMed	for					Go	Clear
<input checked="" type="checkbox"/> Limits		Preview/Index		History	Clipboard		

Display	Abstract	<input type="checkbox"/>	Save	Text	Order	Add to Clipboard
---------	----------	--------------------------	------	------	-------	------------------

Entrez PubMed

☐ 1: *Wien Med Wochenschr* 1999;149(14-15):438-41 Related Articles, Books, LinkOutPubMed
Services**[Immunological mechanisms in allergen-specific immunotherapy:
impact on future trends in type-1 allergy management].**

[Article in German]

Ebner C

Institut für Allgemeine und Experimentelle Pathologie, Wien.

Related
Resources

Specific immunotherapy (SIT) is the only treatment of Type-I allergy that leads to a modulation of the immune response to the eliciting allergen. The repeated administration of high doses of antigen induces a state of "antigen-specific non-responsiveness", i.e. immunologic tolerance to the injected antigen. Accordingly, during and after SIT the proliferative response of allergen-specific T lymphocytes in response to the administered antigen are significantly reduced. According to recent publications, this effect is due to the production of the immunosuppressive cytokine interleukin (IL)-10, which is induced during the treatment. On the other hand, a distinct change in the quality of the immune response to the injected allergen can be observed: the production of the IgE-inducing cytokine IL-4 by T helper cells decreases. Moreover, the release of other proinflammatory cytokines and inflammatory mediators is suppressed. Together, these events result in a marked decrease of symptoms during allergen exposure and reduced reactivity during challenge.

Publication Types:

- Review
- Review, tutorial

PMID: 10584289, UI: 20051678

Display	Abstract	<input type="checkbox"/>	Save	Text	Order	Add to Clipboard
---------	----------	--------------------------	------	------	-------	------------------

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services

National Library
of Medicine

PubMed

PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM
Search PubMed	for					Go	Clear
<input checked="" type="checkbox"/> Limits		Preview/Index		History	Clipboard		

Display	Abstract	<input type="checkbox"/>	Save	Text	Order	Add to Clipboard
---------	----------	--------------------------	------	------	-------	------------------

Entrez PubMed

☐ 1: *Int Immunol* 1999 Oct;11(10):1717-24

Related Articles, Books, LinkOut

Full text article at
intimm.oupjournals.orgPubMed
ServicesCorrected and republished article originally printed in *Int Immunol* 1999
Jul;11(7):1131-8**Suppressive versus stimulatory effects of allergen/cholera toxoid (CTB) conjugates depending on the nature of the allergen in a murine model of type I allergy.**Related
Resources**Wiedermann U, Jahn-Schmid B, Lindblad M, Rask C, Holmgren J, Kraft D, Ebner C**Division of Immunopathology, Institute of General and Experimental Pathology,
University of Vienna, Waehringer Guertel 18-20, 1090 Vienna, Austria.

Recent reports have demonstrated that feeding small amounts of antigen conjugated to the B subunit of cholera toxin (CTB) suppress immune responses in experimental models of certain T(h)1-based autoimmune diseases. We have established a model of aerosol sensitization leading to T(h)2-mediated allergic immune responses in BALB/c mice. In the present study two different antigens, the dietary antigen ovalbumin (OVA) and the inhalant allergen Bet v 1 (the major birch pollen allergen), chemically coupled to recombinant CTB were tested for their potential to influence T(h)2-like immune responses. Intranasal administration of OVA-CTB prior to sensitization with OVA led to a significant decrease of antigen-specific IgE antibody levels, but a marked increase of OVA-specific IgG2a antibodies as compared to non-pretreated, sensitized animals. Antigen-specific lympho-proliferative responses in vitro were reduced by 65% in the pretreated group; IL-5 and IL-4 production were decreased in responder cells of lungs and spleens of nasally pretreated mice. In contrast, mucosal administration of rBet v 1-CTB conjugates prior to sensitization led to an up-regulation of allergen-specific IgE, IgG1 and IgG2a, increased in vitro lympho-proliferative responses as well as augmented production of IL-5, IL-4, IL-10 and IFN-gamma. Intranasal administration prior to sensitization of unconjugated allergens showed also contrasting effects: OVA could not significantly influence antigen-specific antibody or cytokine production, whereas intranasal pretreatment with unconjugated Bet v 1 suppressed allergen-specific immune responses in vivo and in vitro. These results

demonstrated that the two antigens-in conjugated as in unconjugated form-had different effects on the T(h)2 immune responses. We therefore conclude that the tolerogenic or immunogenic properties of CTB-and probably also other antigen-delivery systems-strongly depend on the nature of the coupled antigen-allergen.

Publication Types:

- Corrected and republished article

PMID: 10508191, UI: 99437895

Display	Abstract	<input type="button" value="v"/>	Save	Text	Order	Add to Clipboard
---------	----------	----------------------------------	------	------	-------	------------------


[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)

antigens--in conjugated as in unconjugated form--had different effects on the Th2 immune responses. We therefore conclude that the tolerogenic or immunogenic properties of CTB--and probably also other antigen-delivery systems--strongly depend on the nature of the coupled antigen-allergen.

PMID: 10383946, UI: 99315317

Display	Abstract	<input type="button" value="v"/>	Save	Text	Order	Add to Clipboard
---------	----------	----------------------------------	------	------	-------	------------------

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)

National Library
of Medicine

PubMed

PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM
Search PubMed	▼ for					Go	Clear
Limits		Preview/Index		History	Clipboard		

Entrez PubMed

☐ 1: *J Immunol* 1984 Oct;133(4):1818-24[Related Articles, Books](#)PubMed
Services**Cholera toxin B subunit as a carrier protein to stimulate a mucosal immune response.****McKenzie SJ, Halsey JF**Related
Resources

Horseradish peroxidase (HRP) was covalently coupled to the binding subunit of cholera toxin (CTB) via a two-step glutaraldehyde procedure. The HRP-CTB conjugate was characterized by physiochemical as well as immunochemical methods. Mice were immunized intraduodenally with the HRP-CTB conjugate, with HRP alone, or with a mixture of uncoupled CTB and HRP. The functionally active dose of CTB was 50 micrograms and the HRP dose was in the 30- to 90-micrograms range. Both IgA and IgG antibody responses were measured in serum, intestinal washes, and bile by using a solid phase immunoradiometric assay. Mice immunized with the HRP-CTB conjugate showed a significantly higher level of IgA anti-HRP in intestinal washes and bile, as well as increased levels of serum IgG anti-HRP. Animals that received only HRP or the mixture of CTB and HRP had reduced levels of HRP-specific antibody of either class in both gut washes and bile. The IgA anti-HRP responses in the gut washes were 33- to 120-fold higher when the conjugate was used as the immunogen in comparison with immunization with the CTB + HRP or the HRP alone. Vaccines to stimulate mucosal immunity to any antigenic determinant might thus be prepared by covalent conjugation to effective mucosal immunogens such as CTB.

PMID: 6470484, UI: 84290652

Display	Abstract	▼	Save	Text	Order	Add to Clipboard
---------	----------	---	------	------	-------	------------------

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)